

LOCAL PROJECTION TYPE STABILISATION APPLIED TO INF-SUP STABLE DISCRETISATIONS OF THE OSEEN PROBLEM. PART II

Gunar Matthies^{1,*} and Lutz Tobiska²

¹ Universität Kassel, FB 10, Institut für Mathematik, Heinrich-Plett-Straße 40, 34132 Kassel,
matthies@mathematik.uni-kassel.de

² Otto-von-Guericke-Universität Magdeburg, Institut für Analysis und Numerik, PF 4120,
39014 Magdeburg, tobiska@ovgu.de

Key words: *Stabilised finite elements, Navier–Stokes equations, inf-sup stable elements, local projection*

We consider the Oseen problem

$$\begin{aligned} -\nu\Delta u + (b \cdot \nabla)u + \sigma u + \nabla p &= f && \text{in } \Omega, \\ \operatorname{div} u &= 0 && \text{in } \Omega, \\ u &= 0 && \text{on } \partial\Omega, \end{aligned}$$

where $\nu > 0$ and $\sigma \geq 0$ are constants and $b \in (W^{1,\infty}(\Omega))^d$ with $\operatorname{div} b = 0$ is a given velocity field. The Oseen problem can be considered as a linearisation of the steady ($\sigma = 0$) and the time-discretised non-steady ($\sigma > 0$) Navier–Stokes equations, respectively. In the latter case, one typically has $\sigma \sim 1/\Delta t$ with the time step length Δt of an implicit time integration scheme while the stationary case corresponds to $\Delta t = \infty$.

Inf-sup stable finite element discretisations of the Oseen problem are considered. Hence, no additional pressure stabilisation is needed. However, the standard Galerkin method still suffers in general from global spurious oscillations in the velocity field which are caused by the dominating convection.

Local projection stabilisation methods will be used to overcome this difficulty. The stabilisation is based on a projection from the underlying approximation space onto a discontinuous projection space. Stabilisation is derived from additional weighted L^2 -control on the fluctuation of the gradient of the velocity or only parts of it like divergence and/or derivative in streamline direction.

The convergence analysis for both the one-level and the two-level local projection stabilisation applied to inf-sup stable discretisations of the Oseen problem will be presented in a unified framework.

Two different stabilisation terms

$$S_h^a(u, v) := \sum_{M \in \mathcal{M}_h} \left(\tau_M (\kappa_M^1 (b_M \cdot \nabla) u, \kappa_M^1 (b_M \cdot \nabla) v)_M + \gamma_M (\kappa_M^2 \operatorname{div} u, \kappa_M^2 \operatorname{div} v)_M \right)$$

and

$$S_h^b(u, v) := \sum_{M \in \mathcal{M}_h} \mu_M (\kappa_M^3 \nabla u, \kappa_M^3 \nabla v)_M$$

with non-negative coefficients τ_M, γ_M, μ_M are considered where $\kappa_M^i := \operatorname{id} - \pi_M^i$, $i = 1, 2, 3$, denote the fluctuation operators, $\pi_M^i : L^2(M) \rightarrow D^i(M)$ the local projection operators into finite dimensional spaces. Note that S_h^a introduces control over the fluctuations of the derivative in streamline direction and over the fluctuations of the divergence separately whereas S_h^b controls the fluctuations of the gradient.

We will propose new inf-sup stable pairs of finite element spaces which approximate both velocity and pressure by elements of order r . In contrast to the classical equal order interpolation, the velocity components and the pressure are discretised by different elements. We will show the discrete inf-sup condition for these pairs of finite element spaces. For the case of small viscosity, a uniform error estimate of order $r + 1/2$ will be proved. In the case of discontinuous pressure approximations, an additional term controlling the jumps of the pressure across inner cell faces becomes necessary.

Numerical tests which confirm the theoretical results will be given.

REFERENCES

- [1] G. Matthies and L. Tobiska. Local projection type stabilisation applied to inf-sup stable discretisations of the Oseen problem. accepted for *IMA J. Numer. Anal.*, 2013.
- [2] H.-G. Roos, M. Stynes, and L. Tobiska. Robust numerical methods for singularly perturbed differential equations. Convection-diffusion-reaction and flow problems. Second Edition, Springer-Verlag, Berlin, 2008.