

A SEMI-IMPLICIT METHOD FOR THROMBUS FORMATION IN HAEMODYNAMIC FLUID–STRUCTURE INTERACTION

Richard Schussnig^{1,2,*}, Simon Drey mann³, Alireza Jafarinia^{2,3},
Thomas Hochrainer^{2,3} and Thomas-Peter Fries^{1,2}

¹ Institute of Structural Analysis, Graz University of Technology,
Lessingstraße 25/II, 8010 Graz, Austria, ifb.tugraz.at

² Graz Center of Computational Engineering, Graz University of Technology,
Krenngasse 37/I, 8010 Graz, Austria, gcce.tugraz.at

³ Institute of Strength of Materials, Graz University of Technology,
Kopernikusgasse 24/I, 8010 Graz, Austria, ifl.tugraz.at

Keywords: *Thrombus formation, Fluid–Structure Interaction, Haemodynamics, Incompressible Flow*

Blood clotting and thrombus formation play a key role in many cardiovascular diseases, having a drastic impact on surgical planning and options, risk factors and possible long-term treatment and surveillance. Due to vast developments in the field of computational biomechanics, modern simulation tools can significantly aid clinical decision-making, inspire identifying possible alternatives or highlight risks associated with individual interventions. However, despite these advances, computational modeling of thrombus formation has remained a challenging application, since it combines demanding fluid flow or even fluid-structure interaction problems with complex bio-chemical systems.

In this contribution, we therefore extend a split-step scheme for incompressible non-Newtonian fluid–structure interaction [1] by the thrombus formation model of Menichini et al. [2]. A semi-implicit framework allows coupling only the hyperelastic bulk solid’s displacement and the fluid pressure iteratively, while the equations governing constituent transport and all other involved fields are solved merely once per timestep. Herein, first results in a clinically relevant setting are presented to highlight the efficiency and applicability of the framework.

REFERENCES

- [1] R. Schussnig, D.R.Q. Pacheco, T.-P. Fries, Efficient split-step schemes for fluid-structure interaction involving incompressible generalised Newtonian flows, *Comput Struct*, in press.
- [2] C. Menichini, X.Y. Xu, Mathematical modeling of thrombus formation in idealized models of aortic dissection: initial findings and potential applications. *J Math Biol*, Vol. **73**, pp. 12051226, 2016.